

## APPENDIX

### UNITS, CONSTANTS, AND CONVERSION FACTORS

#### THE INTERNATIONAL SYSTEM OF UNITS (SI)<sup>1</sup>

Table A-1. SI base units.

Name	Symbol	Definition
meter	m	"The meter is the length equal to 1 650 763.73 wavelengths in vacuum of the radiation corresponding to the transition between the levels $2p_{10}$ and $5d_5$ of the krypton-86 atom."
kilogram	kg	"The kilogram is the unit of mass; it is equal to the mass of the international prototype of the kilogram [a cylinder of platinum-iridium]."
second	s	"The second is the duration of 9 192 631 770 periods of the radiation corresponding to the transition between the two hyperfine levels of the ground state of the cesium-133 atom."
ampere	A	"The ampere is the constant current which, if maintained in two straight parallel conductors of infinite length, of negligible circular cross section, and placed 1 meter apart in vacuum, would produce between these conductors a force equal to $2 \times 10^{-7}$ newton per meter of length."
kelvin	K	"The kelvin is the fraction 1/273.16 of the thermodynamic temperature of the triple point of water." The "degree Celsius" is defined by the equation $t = T - T_0$ , where $T$ is the thermodynamic temperature in kelvins and $T_0 = 273.15$ K.
mole	mol	"The mole is the amount of substance of a system which contains as many elementary entities as there are atoms in 0.012 kilogram of carbon 12."
candela	cd	"The candela is the luminous intensity, in a given direction, of a source that emits monochromatic radiation of frequency $540 \times 10^{12}$ hertz and that has a radiant intensity in that direction of 1/683 watt per steradian."

Table A-2. SI derived units.

Name	Symbol (dimensions)	Definition	Value in cgs units <sup>a</sup>
Absorbed dose	Gy (m·s <sup>-2</sup> )	The <i>gray</i> is the absorbed dose when the energy per unit mass imparted to matter by ionizing radiation is one joule per kilogram. (The gray is also used for the ionizing radiation quantities: specific energy imparted, kerma, and absorbed dose index, which have the SI unit joule per kilogram.) 1 rad = $10^{-2}$ Gy.	$10^4$ ergs/gm
Activity	Bq (s <sup>-1</sup> )	The <i>bequerel</i> is the activity of a radionuclide decaying at the rate of one spontaneous nuclear transition per second. 1 Ci (curie) = $3.7 \times 10^{10}$ Bq.	1 sec <sup>-1</sup>
Dose equivalent	Sv (m <sup>2</sup> ·s <sup>-2</sup> )	The <i>sievert</i> is the dose equivalent when the absorbed dose of ionizing radiation multiplied by the dimensionless factors <i>Q</i> (quality factor) and <i>N</i> (product of any other multiplying factors) stipulated by the International Commission on Radiological Protection is one joule per kilogram.	$10^4$ ergs/gm
Electric capacitance	F (m <sup>-2</sup> ·kg <sup>-1</sup> ·s <sup>4</sup> ·A <sup>2</sup> )	The <i>farad</i> is the capacitance of a capacitor between the plates of which there appears a difference of potential of one volt when it is charged by a quantity of electricity equal to one coulomb.	$8.988 \times 10^{11}$ esu
Electric conductance	S (m <sup>-2</sup> ·kg <sup>-1</sup> ·s <sup>3</sup> ·A <sup>2</sup> )	The <i>siemens</i> is the electric conductance of a conductor in which a current of one ampere is produced by an electric potential difference of one volt.	$8.988 \times 10^{11}$ esu (cm/sec)

1. From *Physics Vade Mecum*, edited by H.L. Anderson, American Institute of Physics, New York, 1981.

## APPENDIX

Table A-2 (*continued*)

Name	Symbol (dimensions)	Definition	Value in cgs units <sup>a</sup>
Electric inductance	H (m <sup>2</sup> ·kg·s <sup>-2</sup> ·A <sup>-2</sup> )	The <i>henry</i> is the inductance of a closed circuit in which an electromotive force of one volt is produced when the electric current in the circuit varies uniformly at a rate of one ampere per second.	10 <sup>9</sup> esu (cm)
Electric potential difference, electromotive	V (m <sup>2</sup> ·kg·s <sup>-2</sup> ·A <sup>-1</sup> )	The <i>volt</i> (unit of electric potential difference and electromotive force) is the difference of electric potential between two points of a conductor carrying a constant current of one ampere, when the power dissipated between these points is equal to one watt.	(1/2.998) × 10 <sup>-2</sup> esu (cm <sup>1/2</sup> ·gm <sup>1/2</sup> ·sec <sup>-1</sup> )
Electric resistance	$\Omega$ (m <sup>2</sup> ·kg·s <sup>-3</sup> ·A <sup>-2</sup> )	The <i>ohm</i> is the electric resistance between two points of a conductor when a constant difference of potential of one volt, applied between these two points, produces in this conductor a current of one ampere, this conductor not being the source of any electromotive force.	(1/8.988) × 10 <sup>-11</sup> esu (cm <sup>-1</sup> ·sec)
Energy	J (m <sup>2</sup> ·kg·s <sup>-2</sup> )	The <i>joule</i> is the work done when the point of application of a force of one newton is displaced a distance of one meter in the direction of the force.	10 <sup>7</sup> ergs (cm <sup>2</sup> ·gm·sec <sup>-2</sup> )
Force	N (m·kg·s <sup>-2</sup> )	The <i>newton</i> is that force which, when applied to a body having a mass of one kilogram, gives it an acceleration of one meter per second squared.	10 <sup>5</sup> dyn (cm·gm·sec <sup>-2</sup> )
Frequency	Hz (s <sup>-1</sup> )	The <i>hertz</i> is the frequency of a periodic phenomenon of which the period is one second.	cycles/sec (sec <sup>-1</sup> )
Illuminance	lx (cd·sr·m <sup>-2</sup> )	The <i>lux</i> is the illuminance produced by a luminous flux of one lumen uniformly distributed over a surface of one square meter.	
Luminous flux	lm (cd·sr)	The <i>lumen</i> is the luminous flux emitted in a solid angle of one steradian by a point source having a uniform intensity of one candela.	
Magnetic flux	Wb (m <sup>2</sup> ·kg·s <sup>-2</sup> ·A <sup>-1</sup> )	The <i>weber</i> is the magnetic flux which, linking a circuit of one turn, produces in it an electromotive force of one volt as it is reduced to zero at a uniform rate in one second.	10 <sup>8</sup> Mx (cm <sup>3/2</sup> ·gm <sup>1/2</sup> ·sec <sup>-1</sup> )
Magnetic flux density	T (kg·s <sup>-2</sup> ·A <sup>-1</sup> )	The <i>tesla</i> is the magnetic flux density given by a magnetic flux of one weber per square meter.	10 <sup>4</sup> Gs (cm <sup>-1/2</sup> ·gm <sup>1/2</sup> ·sec <sup>-1</sup> )
Power	W (m <sup>2</sup> ·kg·s <sup>-3</sup> )	The <i>watt</i> is the power which gives rise to the production of energy at the rate of one joule per second.	10 <sup>7</sup> ergs/sec (cm <sup>2</sup> ·gm·sec <sup>-2</sup> )
Pressure or stress	Pa (m <sup>-1</sup> ·kg·s <sup>-2</sup> )	The <i>pascal</i> is the pressure or stress of one newton per square meter.	10 dyn/cm <sup>2</sup> (cm <sup>-1</sup> ·gm·sec <sup>-2</sup> )
Quantity of electricity	C	The <i>coulomb</i> is the quantity of electricity transported in one second by a current of one ampere.	2.998 × 10 <sup>9</sup> esu (cm <sup>3/2</sup> ·gm <sup>1/2</sup> ·sec <sup>-1</sup> )

<sup>a</sup>For more precise work use 2.997 924 58 for 2.998 and 8.987 551 79 for 8.988.

Table A-3. SI prefixes.

Factor	Prefix	Symbol	Factor	Prefix	Symbol
10 <sup>18</sup>	exa	E	10 <sup>-1</sup>	deci	d
10 <sup>15</sup>	peta	P	10 <sup>-2</sup>	centi	c
10 <sup>12</sup>	tera	T	10 <sup>-3</sup>	milli	m
10 <sup>9</sup>	giga	G	10 <sup>-6</sup>	micro	$\mu$
10 <sup>6</sup>	mega	M	10 <sup>-9</sup>	nano	n
10 <sup>3</sup>	kilo	k	10 <sup>-12</sup>	pico	p
10 <sup>2</sup>	hecto	h	10 <sup>-15</sup>	femto	f
10 <sup>1</sup>	deka	da	10 <sup>-18</sup>	atto	a

# UNITS, CONSTANTS, AND CONVERSION FACTORS

Table A-4. Conversion to SI units.

## 1. Acceleration

The gal is a special unit employed in geodesy and geophysics to express the acceleration due to gravity.

1 ft/s <sup>2</sup>	= 0.30480* m/s <sup>2</sup>	1 gal	= 0.01000* m/s <sup>2</sup>
Standard gravity ( <i>g</i> )	= 9.8067 m/s <sup>2</sup>	Sun's surface	= 274.0 m/s <sup>2</sup>

## 2. Angle

1 degree (°)	= $1.7453 \times 10^{-2}$ rad	1 second ("")	= $4.8481 \times 10^{-6}$ rad
1 minute ('')	= $2.9089 \times 10^{-4}$ rad		

## 3. Area

The darcy is a unit for measuring the permeability of porous solids.

1 acre	= 4046.9 m <sup>2</sup>	1 in. <sup>2</sup>	= $6.4516^* \times 10^{-4}$ m <sup>2</sup>
1 are	= 100.00* m <sup>2</sup>	1 square mile (international)	= $2.5900 \times 10^6$ m <sup>2</sup>
1 barn (b)	= $1.0000^* \times 10^{-28}$ m <sup>2</sup>	1 square mile (statute) <sup>†</sup>	= $2.5900 \times 10^6$ m <sup>2</sup>
1 circular mil	= $5.0671 \times 10^{-10}$ m <sup>2</sup>	1 square (building)	= 9.2903 m <sup>2</sup>
1 darcy	= $9.8692 \times 10^{-13}$ m <sup>2</sup>	1 square rod (rd <sup>2</sup> ), square pole, or square perch	= 25.293 m <sup>2</sup>
1 ft <sup>2</sup>	= 0.092 903 m <sup>2</sup>	1 square yard (yd <sup>2</sup> )	= 0.83613 m <sup>2</sup>
1 hectare	= 10 000* m <sup>2</sup>		

## 4. Density

1 grain/gal (U.S. liquid)	= 0.017118 kg/m <sup>3</sup>	1 ton (short)/yd <sup>3</sup>	= 1186.6 kg/m <sup>3</sup>
1 oz (avoirdupois)/in. <sup>3</sup>	= 1730.0 kg/m <sup>3</sup>	Density of water (4°C)	= 999.97 kg/m <sup>3</sup>
1 lb/ft <sup>3</sup>	= 16.018 kg/m <sup>3</sup>	Density of mercury (0°C)	= 13595 kg/m <sup>3</sup>
1 lb/in. <sup>3</sup>	= 27680 kg/m <sup>3</sup>	Solar mass/cubic parsec	= $6.770 \times 10^{-20}$ kg/m <sup>3</sup>
1 lb/gal (U.S. liquid)	= 119.83 kg/m <sup>3</sup>	STP gas density for molecular weight <i>M</i> <sub>0</sub>	= 0.044615 <i>M</i> <sub>0</sub> kg/m <sup>3</sup>
1 ton (long)/yd <sup>3</sup>	= 1328.9 kg/m <sup>3</sup>		

## 5. Electricity and magnetism

A = ampere, C = coulomb, F = farad, H = henry, Ω = ohm, S = siemens, V = volt, T = tesla, Wb = weber, \* = exact value.

1 abampere	= 10.000* A	1 ohm centimeter	= $1.0000^* \times 10^{-2}$ Ω·m
1 abcoulomb	= 10.000* C	1 ohm circular-mil per foot	= $1.6624 \times 10^{-9}$ Ω·m
1 abfarad	= $1.0000^* \times 10^9$ F	1 statampere	= $3.3356 \times 10^{-10}$ A
1 abhenry	= $1.0000^* \times 10^{-9}$ H	1 statcoulomb	= $3.3356 \times 10^{-10}$ C
1 abmho	= $1.0000^* \times 10^9$ S	1 statfarad	= $1.1127 \times 10^{-12}$ F
1 abohm	= $1.0000^* \times 10^{-9}$ Ω	1 stathenry	= $8.9876 \times 10^{11}$ H
1 abvolt	= $1.0000^* \times 10^{-8}$ V	1 statmho	= $1.1127 \times 10^{-12}$ S
1 ampere hour	= 3600* C	1 statohm	= $8.9876 \times 10^{11}$ Ω
1 emu of capacitance	= $1.0000^* \times 10^8$ F	1 statvolt	= 299.79 V
1 emu of current	= 10.000* A	1 unit pole	= $1.2566 \times 10^{-7}$ Wb
1 emu of electric potential	= $1.0000^* \times 10^{-8}$ V	Potential of electron at 1st Bohr orbit	= 27.212 V
1 emu of inductance	= $1.0000^* \times 10^{-9}$ H	Ionization potential from 1st Bohr orbit	= 13.606 V
1 emu of resistance	= $1.0000^* \times 10^{-9}$ Ω	Nuclear electric field at 1st Bohr orbit	= $5.140 \times 10^{11}$ V/m
1 esu of capacitance	= $1.1127 \times 10^{-12}$ F	Current in 1st Bohr orbit	= $1.054 \times 10^{-3}$ A
1 esu of current	= $3.3356 \times 10^{-10}$ A	Dipole moment of nucleus and electron in 1st Bohr orbit	= $0.8478 \times 10^{-29}$ C·m
1 esu of electric potential	= $2.9979 \times 10^2$ V	Magnetic field, atomic unit	= 1715 T
1 esu of inductance	= $8.9876 \times 10^{11}$ H	Field at nucleus due to electron in 1st Bohr orbit	= $9.9551 \times 10^6$ A/m
1 esu of resistance	= $8.9876 \times 10^{11}$ Ω	Magnetic moment, atomic unit	= $2.542 \times 10^{-21}$ J/T
1 faraday (based on <sup>12</sup> C)	= $9.6487 \times 10^4$ C	Earth magnetic moment	= $7.98 \times 10^{22}$ J/T
1 faraday (chemical)	= $9.6496 \times 10^4$ C		
1 faraday (physical)	= $9.6522 \times 10^4$ C		
1 gamma	= $1.0000^* \times 10^{-9}$ T		
1 gauss	= $1.0000^* \times 10^{-4}$ T		
1 gilbert	= $7.9577 \times 10^{-1}$ A (amp. turns)		
1 maxwell	= $1.0000^* \times 10^{-8}$ Wb		
1 mho	= 1.0000* S		
1 oersted	= 79.577 A/m		

## 6. Energy

Btu = British thermal unit (thermochemical), 1 Btu (International Table) = 1.000 67 Btu (thermochemical); cal = calorie (thermochemical), 1 cal (International Table) = 1.000 67 cal (thermochemical); J = joule; W = watt.

1 Btu	= 1054.4 J	1 foot-poundal	= 0.042140 J
1 Btu (mean)	= 1055.9 J	1 kilowatt hour (kW·h)	= $3.6000^* \times 10^6$ J
1 Btu (39°F)	= 1059.7 J	1 therm	= $1.0551 \times 10^8$ J

## APPENDIX

Table A-4 (*continued*)

1 Btu (60°F)	= 1054.7 J	1 ton (nuclear equivalent of TNT)	= $4.184 \times 10^9$ J
1 calorie	= 4.1840* J	1 watt hour (W·h)	= 3600* J
1 calorie (mean)	= 4.1900 J	1 watt second (W·s)	= 1.0000* J
1 calorie (15°C)	= 4.1858 J	Energy of unit wave number ( $hc$ )	= $1.9865 \times 10^{-25}$ J
1 calorie (20°C)	= 4.1819 J	Mass energy of unit atomic weight	= $1.4924 \times 10^{-10}$ J
1 kilocalorie	= 4184.0* J		
1 electron volt (eV)	= $1.6022 \times 10^{-19}$ J		
1 erg	= $1.0000 \times 10^{-7}$ J		
1 foot-pound (ft-lbf)	= 1.3558 J		
Note: 1 quad = $10^{15}$ Btu			
1 quad per year = 0.472 million barrels of oil per day (1 barrel = 42 gallons)			
= 1 trillion cubic feet of gas per year			
= 44.4 million tons of coal per year (for medium heating value coal at 22.5 Btu/ton)			
= 33.4 million kilowatts of electricity			
= 293 billion kilowatt-hours of electricity per year at 100% efficiency			
= 95.2 billion kilowatt-hours of electricity per year at 32.5% efficiency			
See Chaps. 10.00 (Energy Demand) and 11.00 (Energy Supply).			
<i>7. Force</i>			
1 dyne	= $1.0000 \times 10^{-5}$ N	1 lbf/lb (thrust/weight [mass] ratio)	= 9.8067 N
1 kilogram-force	= 9.8067 N	1 poundal	= 0.13825 N
1 kip (1000 lbf)	= 4448.2 N	1 ton-force (2000 lbf)	= 8896.4 N
1 ounce-force	= 0.27801 N	Proton-electron attraction at distance $a_0$	= $8.238 \times 10^{-8}$ N
1 pound-force (lbf)	= 4.4482 N		
<i>8. Frequency</i>			
1 hertz (Hz)	= 1 cycle/s	Frequency of free electron in magnetic field $H$	= $2.7993 \times 10^{10} H$ Hz·T <sup>-1</sup>
1 kayser	= $3 \times 10^{10}$ Hz		
Rydberg frequency ( $cR_\infty$ )	= $3.2898 \times 10^{15}$ Hz	Plasma frequency associated with electron density $N_e$	= $8.979 N_e^{1/2}$ Hz ( $N_e$ in m <sup>-3</sup> )
Frequency of 1st Bohr orbit ( $2cR_\infty$ )	= $6.5797 \times 10^{15}$ Hz		
<i>9. Heat</i>			
Btu = British thermal unit (thermochemical), 1 Btu (International Table) = 1.000 67 Btu (thermochemical); cal = calorie (thermochemical), 1 cal (International Table) = 1.000 67 cal (thermochemical); J = joule, K = kelvin; W = watt; h = hour.			
Thermal conductivity $k$ :			
1 Btu·ft/h·ft <sup>2</sup> ·°F	= 1.7296 W/m·K	1 Btu·in./s·ft <sup>2</sup> ·°F	= 518.87 W/m·K
Thermal conductance $C$ :			
1 Btu/h·ft <sup>2</sup> ·°F	= 5.6745 W/m <sup>2</sup> ·K		
Heat capacity:			
1 Btu/lb·°F	= 4184.0* J/kg·K	1 cal/s	= 4.1840* W
1 cal/g·°C	= 4184.0* J/kg·K		
Thermal resistance $R$ :			
1 °F·h·ft <sup>2</sup> /Btu	= 0.176 23 K·m <sup>2</sup> /W	Thermal resistivity:	
1 clo	= 0.20037 K·m <sup>2</sup> /W	1 °F·h·ft <sup>2</sup> /Btu·in.	= 6.9381 K·m/W
Thermal diffusivity:			
1 ft <sup>2</sup> /h	= $2.5806 \times 10^{-5}$ m <sup>2</sup> /s		
<i>10. Length</i>			
1 angstrom (Å)	= $1.0000 \times 10^{-10}$ m	1 microinch	= $2.5400 \times 10^{-8}$ m
1 atomic unit ( $a_0$ )	= $0.52918 \times 10^{-10}$ m	1 micron ( $\mu\text{m}$ )	= $1.0000 \times 10^{-6}$ m
1 astronomical unit (AU)	= $1.4960 \times 10^{11}$ m	1 mil	= $2.5400 \times 10^{-5}$ m
1 cable's length	= 219 m	1 mile (int. nautical)	= 1852.0* m
1 chain	= 20.117 m	1 mile (U.S. nautical)	= 1852.0* m
1 electron radius ( $r_e$ )	= $2.818 \times 10^{-15}$ m	1 mile (international)	= 1609.3 m
1 fathom	= 1.8288 m	1 mile (U.S. statute)	= 1609.3 m
1 fermi (femtometer) (fm)	= $1.0000 \times 10^{-15}$ m	1 parsec (pc)	= $3.0857 \times 10^{16}$ m
1 foot (ft)	= 0.30480* m	1 pica (printer's)	= $4.2175 \times 10^{-3}$ m
1 foot (U.S. survey)	= 0.30480 m	1 point (printer's)	= $3.5146 \times 10^{-4}$ m
1 furlong	= 201.17 m	1 rod	= 5.0292 m
1 hand	= 0.10160 m	1 solar radius ( $R_{\odot}$ )	= $6.960 \times 10^8$ m
1 inch (in.)	= 0.02540* m	Wavelength of 1-eV photon ( $hc/eV$ )	= $1.2399 \times 10^{-6}$ m
1 league (land)	= 4828 m		

# UNITS, CONSTANTS, AND CONVERSION FACTORS

Table A-4 (*continued*)

1 light year (ly)	$= 9.4606 \times 10^{15}$ m	1 x unit	$= 1.002 \times 10^{-13}$ m
1 link (Gunther's or surveyors')	$= 0.20117$ m	1 yard	$= 0.91440^*$ m
<b>11. Light</b>			
cd = candela, lm = lumen, [1 lumen = flux from $(1/60\pi)$ cm <sup>2</sup> of blackbody at 2044K], lx = lux.			
1 apostilb	$= 1 \text{ lm/m}^2$ for perfectly diffusing surface	1 lambert	$= 3183.1 \text{ cd/m}^2$
1 cd/in. <sup>2</sup>	$= 1550.0 \text{ cd/m}^2$	1 lumen of maximum-visibility radiation (5550 Å)	$= 1.470 \times 10^{-3}$ W
1 foot-candle	$= 10.764$ lx	1 stilb (sb)	$= 10\,000^* \text{ cd/m}^2$
1 foot-lambert	$= 3.4263 \text{ cd/m}^2$	1 phot	$= 10\,000^* \text{ lx}$
<b>12. Mass</b>			
1 atomic unit (electron) ( $m_e$ )	$= 9.1095 \times 10^{-34}$ kg	1 ounce (avoirdupois)	$= 2.8350 \times 10^{-2}$ kg
1 atomic mass unit ( <sup>12</sup> C scale) (amu)	$= 1.66057 \times 10^{-27}$ kg	1 ounce (troy or apothecary)	$= 3.1103 \times 10^{-2}$ kg
1 carat (metric)	$= 2.0000^* \times 10^{-4}$ kg	1 pennyweight (troy)	$= 1.5552 \times 10^{-3}$ kg
1 dram, apothecary	$= 3.8879 \times 10^{-3}$ kg	1 pound (lb avoirdupois)	$= 0.45359$ kg
1 dram, avoirdupois	$= 1.7718 \times 10^{-3}$ kg	1 pound (troy or apothecary)	$= 0.37324$ kg
1 gamma	$= 1.0000^* \times 10^{-9}$ kg	1 quintal (q)	$= 100.00$ kg
1 grain	$= 6.4799 \times 10^{-6}$ kg	1 scruple	$= 1.2960 \times 10^{-3}$ kg
1 hundredweight (gross or long)	$= 50.802$ kg	1 slug	$= 14.594$ kg
1 hundred weight (net or short)	$= 45.359$ kg	1 ton (assay)	$= 0.029167$ kg
1 kgf·s <sup>2</sup> /m	$= 9.8067$ kg	1 ton (long, 2240 lb)	$= 1016.0$ kg
1 ton (short, 2000 lb)		1 ton (short, 2000 lb)	$= 907.18$ kg
1 ton (metric ton)		1 ton (metric ton)	$= 1.0000^*$ kg
<b>13. Mass per unit length</b>			
1 denier	$= 1.1111 \times 10^{-7}$ kg/m	1 tex	$= 1.0000^* \times 10^{-6}$ kg/m
<b>14. Mass per unit time</b>			
1 perm (0°C)	$= 5.7214 \times 10^{-11}$ kg/Pa·s·m <sup>2</sup>	1 lb/hp·h	$= 1.6897 \times 10^{-7}$ kg/J
1 perm-in. (0°C)	$= 1.4532 \times 10^{-12}$ kg/Pa·s·m	1 ton (short)/h	$= 0.25200$ kg/s
1 lb/h	$= 1.2600 \times 10^{-4}$ kg/s		
<b>15. Power</b>			
1 Btu (int.)/h	$= 0.29307$ W	1 horsepower (boiler)	$= 9809.5$ W
1 Btu (int.)/s	$= 1055.1$ W	1 horsepower (electric)	$= 746.00^*$ W
1 Btu (thermochem.)/h	$= 0.29288$ W	1 horsepower (metric)	$= 735.50$ W
1 cal (thermochem.)/s	$= 4.1840^*$ W	1 horsepower (water)	$= 746.04$ W
1 force de cheval	$= 735.5$ W	1 horsepower (U.K.)	$= 745.70$ W
1 erg/s	$= 1.0000^* \times 10^{-7}$ W	1 ton (refrigeration)	$= 3516.8$ W
1 ft-lbf/h	$= 3.7662 \times 10^{-4}$ W	Star, $M_{bol} = 0$ radiation	$= 2.97 \times 10^{28}$ W
1 horsepower (550 ft-lbf/s)	$= 745.70$ W	Solar luminosity	$= 3.826 \times 10^{26}$ W
<b>16. Pressure or stress (force per unit area)</b>			
1 atmosphere (standard)	$= 101\,325^*$ Pa	1 inch of water (39.2°F)	$= 249.08$ Pa
1 atmosphere (technical = 1 kgf/cm <sup>2</sup> )	$= 98\,066.5^*$ Pa	1 kgf/cm <sup>2</sup>	$= 98\,066.5^*$ Pa
1 bar	$= 100\,000^*$ Pa	1 kip/in. <sup>2</sup> (ksi)	$= 6.8948 \times 10^6$ Pa
1 cm Hg(0°C)	$= 133.3$ Pa	1 millibar	$= 100.00^*$ Pa
1 centimeter of water (4°C)	$= 98.064$ Pa	1 newton/cm <sup>2</sup>	$= 10000^*$ Pa
1 dyne/cm <sup>2</sup>	$= 0.100\,00^*$ Pa	1 poundal/ft <sup>2</sup>	$= 1.4882$ Pa
1 foot of water (39.2°F)	$= 2989.0$ Pa	1 lbf/ft <sup>2</sup>	$= 47.880$ Pa
1 gf/cm <sup>2</sup>	$= 98.0665^*$ Pa	1 lbf/in. <sup>2</sup> (psi)	$= 6894.8$ Pa
1 inch of mercury (32°F)	$= 3386.4$ Pa	1 torr (mm Hg, 0°C)	$= 133.32$ Pa
<b>17. Temperature</b>			
Degree Celsius	$T_K = T_C + 273.15$	Triple point of natural water	
Degree Fahrenheit	$t_C = (t_F - 32)/1.8$		$= 273.16$ K
Degree Fahrenheit	$T_K = (t_F + 459.67)/1.8$	Elementary temperature ( $\partial h/r_e k$ )	$= 8.1262 \times 10^{11}$ K
Degree Rankine	$T_K = T_R/1.8$	Temperature of 1 eV	$= 11\,605$ K
Kelvin	$t_C = T_K - 273.15$		
<b>18. Time</b>			
1 day	$= 86400^*$ s	1 year (sidereal)	$= 3.1558 \times 10^7$ s
1 day (sidereal)	$= 86164$ s	1 year (tropical)	$= 3.1557 \times 10^7$ s
1 hour	$= 3600.0^*$ s	1 atomic second ( $s_A$ )	$= 9192631770$
1 hour (sidereal)	$= 3590.2$ s		<sup>133</sup> Cs cycles

## APPENDIX

Table A-4 (*continued*)

1 minute	$= 60.000^* \text{ s}$	1 atomic unit (1st Bohr orbit/ $2\pi$ ) ( $\tau_0$ )	$= 2.4189 \times 10^{-17} \text{ s}$
1 second (sidereal)	$= 0.99727 \text{ s}$	Jordan's elementary time ( $r_e/c$ )	$= 9.3996 \times 10^{-24} \text{ s}$
1 year (365 days)	$= 3.1536 \times 10^7 \text{ s}$		
<b>19. Torque</b>			
1 dyne·cm	$= 1.0000^* \times 10^{-7} \text{ N}\cdot\text{m}$	1 lbf·in.	$= 0.11298 \text{ N}\cdot\text{m}$
1 kgf·m	$= 9.8067 \text{ N}\cdot\text{m}$	1 lbf·ft	$= 1.3558 \text{ N}\cdot\text{m}$
1 ozf·in.	$= 0.0070616 \text{ N}\cdot\text{m}$		
<b>20. Velocity</b>			
1 ft/s	$= 0.30480^* \text{ m/s}$	1 AU per year	$= 4.7406 \text{ km/s}$
1 in./s	$= 0.02540^* \text{ m/s}$	1 parsec per year	$= 9.7781 \times 10^8 \text{ m/s}$
1 km/h	$= 0.27778 \text{ m/s}$	Electron in Bohr orbit	$= 2.1877 \times 10^6 \text{ m/s}$
1 knot (international)	$= 0.51444 \text{ m/s}$	1-eV electron	$= 5.9309 \times 10^5 \text{ m/s}$
1 mi/h (international)	$= 0.44704^* \text{ m/s}$	Angular velocity of Earth on its axis	$= 7.2921 \times 10^{-5} \text{ rad/s}$
1 mi/s (international)	$= 1609.3 \text{ m/s}$	Mean angular velocity of Earth in its orbit	$= 1.9910 \times 10^{-7} \text{ rad/s}$
1 mi/h (international)	$= 1.6093 \text{ km/h}$		
Velocity of light ( $c$ )	$= 2.9979 \times 10^8 \text{ m/s}$		
<b>21. Viscosity</b>			
1 centipoise	$= 1.0000^* \times 10^{-3} \text{ Pa}\cdot\text{s}$	1 lb/ft·s	$= 1.4882 \text{ Pa}\cdot\text{s}$
1 centistokes	$= 1.0000^* \times 10^{-6} \text{ m}^2/\text{s}$	1 lbf·s/ft <sup>2</sup>	$= 47.880 \text{ Pa}\cdot\text{s}$
1 ft <sup>2</sup> /s	$= 0.092030 \text{ m}^2/\text{s}$	1 lbf·s/in. <sup>2</sup>	$= 6894.8 \text{ Pa}\cdot\text{s}$
1 poise	$= 0.10000^* \text{ Pa}\cdot\text{s}$	1 rhe	$= 10.000^* (\text{Pa}\cdot\text{s})^{-1}$
1 poundal·s/ft <sup>2</sup>	$= 1.4882 \text{ Pa}\cdot\text{s}$	1 slug/ft·s	$= 47.880 \text{ Pa}\cdot\text{s}$
1 lb/ft·h	$= 4.1338 \times 10^{-4} \text{ Pa}\cdot\text{s}$	1 stokes	$= 1.0000^* \times 10^{-4} \text{ m}^2/\text{s}$
<b>22. Volume</b>			
1 acre-foot	$= 1233.5 \text{ m}^3$	1 ft <sup>3</sup>	$= 0.028317 \text{ m}^3$
1 barrel (oil, 42 gal)	$= 0.15899 \text{ m}^3$	1 gallon (Canadian liquid)	$= 4.5461 \times 10^{-3} \text{ m}^3$
1 barrel (bbl), liquid	$= 31 \text{ to } 42 \text{ gallons}$	1 gallon (U.K. liquid)	$= 4.5461 \times 10^{-3} \text{ m}^3$
1 barrel (bbl), standard for fruits, vegetables, and other dry commodities except cranberries	$= 7056 \text{ cubic inches}$ $= 105 \text{ dry quarts}$ $= 3.281 \text{ bushels, struck measure}$ $= 5826 \text{ cubic inches}$ $= 86_{64}^{45} \text{ dry quarts}$ $= 2.709 \text{ bushels, struck measure}$	1 gallon (U.S. dry) 1 gallon (U.S. liquid) 1 gill (U.K.) 1 gill (U.S.) 1 in. <sup>3</sup> 1 liter	$= 4.4049 \times 10^{-3} \text{ m}^3$ $= 3.7854 \times 10^{-3} \text{ m}^3$ $= 1.4207 \times 10^{-4} \text{ m}^3$ $= 1.1829 \times 10^{-4} \text{ m}^3$ $= 1.6387 \times 10^{-5} \text{ m}^3$ $= 1.0000^* \times 10^{-3} \text{ m}^3$
1 barrel (bbl), standard, cranberry	$= 2.3597 \times 10^{-3} \text{ m}^3$	1 ounce (U.K. fluid) 1 ounce (U.S. fluid)	$= 2.8413 \times 10^{-5} \text{ m}^3$ $= 2.9574 \times 10^{-5} \text{ m}^3$
1 board foot	$= 3.5239 \times 10^{-2} \text{ m}^3$	1 cubic parsec	$= 2.9380 \times 10^{49} \text{ m}^3$
1 bushel (U.S.)		1 peck (U.S.)	$= 8.8098 \times 10^{-3} \text{ m}^3$
1 bushel (bu), struck measure (U.S.)	$= 2150.42^* \text{ in.}^3$	1 pint (U.S. dry) 1 pint (U.S. liquid)	$= 5.5061 \times 10^{-4} \text{ m}^3$ $= 4.7318 \times 10^{-4} \text{ m}^3$
1 bushel, heaped (U.S.)	$= 1.278 \text{ bushels, struck measure}$	1 quart (U.S. dry) 1 quart (U.S. liquid)	$= 1.1012 \times 10^{-3} \text{ m}^3$ $= 9.4635 \times 10^{-4} \text{ m}^3$
1 bushel (bu), struck measure (British Imperial)	$= 1.032 \text{ U.S. bushels, struck measure}$	1 stere	$= 1.0000^* \text{ m}^3$
1 cord (cd) (firewood)	$= 128^* \text{ ft}^3$	Solar volume ( $4\pi R_{\odot}^3/3$ )	$= 1.4122 \times 10^{27} \text{ m}^3$
1 cup (measuring)	$= 2.3659 \times 10^{-4} \text{ m}^3$	1 tablespoon	$= 1.4787 \times 10^{-5} \text{ m}^3$
1 dram (U.S. fluid)	$= 3.6967 \times 10^{-6} \text{ m}^3$	1 teaspoon	$= 4.9289 \times 10^{-6} \text{ m}^3$
1 drachm (U.K. fluid)	$= 3.5516 \times 10^{-6} \text{ m}^3$	1 ton (register)	$= 2.8317 \text{ m}^3$
1 fluid ounce (U.S.)	$= 2.9574 \times 10^{-5} \text{ m}^3$	1 yd <sup>3</sup>	$= 0.76455 \text{ m}^2$

# UNITS, CONSTANTS, AND CONVERSION FACTORS

Table A-5. Precise physical constants.<sup>1</sup>

Quantity	Symbol (expression)	Value in SI (cgs) units	Error (ppm)
1. Speed of light in vacuum	$c$	$2.997\ 924\ 58 \times 10^8 \text{ m}\cdot\text{s}^{-1}$ ( $10^{10} \text{ cm}\cdot\text{sec}^{-1}$ )	0.004
2. Elementary charge	$e$	$1.602\ 189\ 2 \times 10^{-19} \text{ C}$ ( $10^{-20} \text{ emu}$ )	2.9
		( $4.803\ 242 \times 10^{-10} \text{ esu}$ )	2.9
3. Planck's constant	$\hbar$	$6.626\ 176 \times 10^{-34} \text{ J}\cdot\text{s}$ ( $10^{-27} \text{ erg}\cdot\text{sec}$ )	5.4
	$\hbar = h/2\pi$	$1.054\ 588\ 7 \times 10^{-34} \text{ J}\cdot\text{s}$ ( $10^{-27} \text{ erg}\cdot\text{sec}$ )	5.4
4. Electron rest mass	$m_e$	$0.910\ 953\ 4 \times 10^{-30} \text{ kg}$ ( $10^{-27} \text{ gm}$ )	5.1
5. Avogadro constant recent value	$N_A$	$6.022045 \times 10^{23} \text{ mol}^{-1}$ ( $10^{23} \text{ mol}^{-1}$ )	5.1
		$6.022\ 097\ 8 \times 10^{23} \text{ mol}^{-1}$ ( $10^{23} \text{ mol}^{-1}$ )	1.0
6. Molar gas constant	$R$	$8.314\ 41 \times 10^0 \text{ J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$ ( $10^7 \text{ erg}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$ )	31
7. Boltzmann constant	$k = R/N_A$	$1.380\ 662 \times 10^{-23} \text{ J}\cdot\text{K}^{-1}$ ( $10^{-16} \text{ erg}\cdot\text{K}^{-1}$ )	32
8. Gravitational constant	$G$	$6.672\ 0 \times 10^{-11} \text{ N}\cdot\text{m}^2\cdot\text{kg}^{-2}$ ( $10^{-8} \text{ dyn}\cdot\text{cm}^2\cdot\text{gm}^{-2}$ )	615
9. Molar volume ( $273.15^\circ\text{K}$ , $p_0 = 1 \text{ atm}$ )	$V_m = RT_0/p_0$	$22.413\ 83 \times 10^{-3} \text{ m}^3\cdot\text{mol}^{-1}$ ( $10^3 \text{ cm}^3\cdot\text{mol}^{-1}$ )	31
10. Faraday constant	$F = N_A e$	$9.648\ 456 \times 10^4 \text{ C}\cdot\text{mol}^{-1}$ ( $10^3 \text{ emu}\cdot\text{mol}^{-1}$ )	2.8
11. Rydberg constant recent value	$R_x = [4\pi\epsilon_0]^{-2}(m_e e^4 / 4\pi\hbar^3 c)$	$1.097\ 373\ 177 \times 10^7 \text{ m}^{-1}$ ( $10^5 \text{ cm}^{-1}$ )	0.07
		$1.097\ 373\ 147\ 6 \times 10^7 \text{ m}^{-1}$ ( $10^5 \text{ cm}^{-1}$ )	0.0003
12. Fine structure constant recent value	$\alpha^{-1} = [4\pi\epsilon_0](\hbar c/e^2)$	137.036 04	0.11
		137.035 963	
13. Classical electron radius	$r_e = [4\pi\epsilon_0]^{-1}(e^2/m_e c^2)$	$2.817\ 938\ 0 \times 10^{-15} \text{ m}$ ( $10^{-13} \text{ cm}$ )	2.5
14. Specific electron charge	$e/m_e$	$1.758\ 804\ 7 \times 10^{11} \text{ C}\cdot\text{kg}^{-1}$ ( $10^7 \text{ emu}\cdot\text{gm}^{-1}$ )	2.8
15. Electron Compton wavelength	$\lambda_c = \hbar/m_e c = \alpha^{-1} r_e$	$3.861\ 590\ 5 \times 10^{-13} \text{ m}$ ( $10^{-11} \text{ cm}$ )	1.6
16. Bohr radius	$a_0 = \alpha^{-2} r_e$	$0.529\ 177\ 06 \times 10^{-10} \text{ m}$ ( $10^{-8} \text{ cm}$ )	0.82
17. Magnetic flux quantum	$\Phi_0 = [c]^{-1}(\hbar c/2e)$	$2.067\ 850\ 6 \times 10^{-16} \text{ T}\cdot\text{m}^2$ ( $10^{-7} \text{ Gs}\cdot\text{cm}^2$ )	2.6
	$h/e$	$4.135\ 701 \times 10^{-15} \text{ J}\cdot\text{s}\cdot\text{C}^{-1}$ ( $10^{-7} \text{ erg}\cdot\text{sec}\cdot\text{emu}^{-1}$ )	2.6
18. Quantum of circulation	$h/2m_e$	$3.636\ 945\ 5 \times 10^{-4} \text{ J}\cdot\text{Hz}^{-1}\cdot\text{kg}^{-1}$ ( $10^0 \text{ erg}\cdot\text{sec}\cdot\text{gm}^{-1}$ )	1.6
19. Atomic mass unit	$1 \text{ u} = \text{gm}\cdot\text{mol}^{-1}/N_A$	$1.660\ 565\ 5 \times 10^{-27} \text{ kg}$ ( $10^{-24} \text{ gm}$ )	5.1
20. Proton rest mass	$m_p$	$1.672\ 648\ 5 \times 10^{-27} \text{ kg}$ ( $10^{-24} \text{ gm}$ )	5.1
		1.007 276 470 u (amu)	0.011
	$m_p/m_e$	1836.151 52	0.38
21. Neutron rest mass	$m_n$	$1.674\ 954\ 3 \times 10^{-27} \text{ kg}$ ( $10^{-24} \text{ gm}$ )	5.1
		1.008 665 012 u (amu)	0.037
22. Electron g factor recent value	$\frac{1}{2}g_e = \mu_e/\mu_B$	1.001 159 656 7	0.0035
		1.001 159 652 200	0.0004
23. Bohr magneton	$\mu_B = [c](e\hbar/2m_e c)$	$9.274\ 078 \times 10^{-24} \text{ J}\cdot\text{T}^{-1}$ ( $10^{-21} \text{ erg}\cdot\text{Gs}^{-1}$ )	3.9
24. Nuclear magneton	$\mu_N = [c](e\hbar/2m_p c)$	$5.050\ 824 \times 10^{-27} \text{ J}\cdot\text{T}^{-1}$ ( $10^{-24} \text{ erg}\cdot\text{Gs}^{-1}$ )	3.9
25. Electron magnetic moment	$\mu_e$	$9.284\ 832 \times 10^{-24} \text{ J}\cdot\text{T}^{-1}$ ( $10^{-21} \text{ erg}\cdot\text{Gs}^{-1}$ )	3.9
26. Proton magnetic moment	$\mu_p$	$1.410\ 617\ 1 \times 10^{-26} \text{ J}\cdot\text{T}^{-1}$ ( $10^{-23} \text{ erg}\cdot\text{Gs}^{-1}$ )	3.9
	$\mu_e/\mu_p$	658.210 688 0	0.010
27. Proton gyromagnetic ratio	$\gamma_p$	$2.675\ 198\ 7 \times 10^8 \text{ rad}\cdot\text{s}^{-1}\cdot\text{T}^{-1}$ ( $10^4 \text{ rad}\cdot\text{sec}^{-1}\cdot\text{Gs}^{-1}$ )	2.8
28. Stefan-Boltzmann constant	$\sigma = (\pi^2/60)k^4/\hbar^3c^2$	$5.670\ 32 \times 10^{-8} \text{ W}\cdot\text{m}^{-2}\cdot\text{K}^{-4}$ ( $10^{-5} \text{ erg}\cdot\text{sec}^{-1}\cdot\text{cm}^{-2}\cdot\text{K}^{-4}$ )	125
29. First radiation constant	$C_1 = 2\pi\hbar c^2$	$3.741\ 832 \times 10^{-16} \text{ W}\cdot\text{m}^2$ ( $10^{-5} \text{ erg}\cdot\text{sec}^{-1}\cdot\text{cm}^2$ )	5.4
30. Second radiation constant	$C_2 = hc/k$	$1.438\ 786 \times 10^{-2} \text{ m}\cdot\text{K}$ ( $10^0 \text{ cm}^{-2}\cdot\text{K}$ )	31

## Energy equivalents

Quantity	Symbol (expression)	Value	Error (ppm)
Atomic mass unit	$u$	931.501 6 MeV	2.8
Proton mass	$m_p$	938.279 6 MeV	2.8
Neutron mass	$m_n$	939.573 1 MeV	2.8
Electron mass	$m_e$	0.511 003 4 MeV	2.8
Electron volt	1 eV/K	11 604.50 K	31
	1 eV/hc	$8\ 065\ 479 \text{ cm}^{-1}$	2.6
	1 eV/h	$2.417\ 969\ 6 \times 10^{14} \text{ Hz}$	2.6
	1 eV	$1.602\ 189\ 2 \times 10^{-12} \text{ ergs}$	2.9
Planck's constant	$\hbar$	$6.582\ 173 \times 10^{-22} \text{ MeV}\cdot\text{sec}$	2.6
	$\hbar c$	$1.973\ 285\ 8 \times 10^{-11} \text{ MeV}\cdot\text{cm}$	2.6
	$(\hbar c)^2$	0.389 385 7 GeV <sup>2</sup> ·mb	5.2
Rydberg constant	$R_x hc$	13.605 804 eV	2.6
Voltage-wavelength product	$V\lambda$	12 398.520 eV·Å	2.6
Gas constant	$R$	1.987 19 cal·mol <sup>-1</sup> ·K <sup>-1</sup>	31

## APPENDIX

Table A-6. Mathematical constants.<sup>1</sup>

Constant	Number	Log
$\pi$	3.141 592 653 6	0.497 149 872 7
$2\pi$	6.283 185 307 2	0.798 179 868 4
$4\pi$	12.566 370 614 4	1.099 209 864 0
$\pi^2$	9.869 604 401 1	0.994 299 745 4
$\sqrt{\pi}$	1.772 453 850 9	0.248 574 936 3
$e$	2.718 281 828 5	0.434 294 481 9
$\text{mod} = M = \log e$	0.434 294 481 9	1.637 784 311 3
$1/M = \ln 10$	2.302 585 093 0	0.362 215 688 7
$2$	2.000 000 000 0	0.301 029 995 7
$\sqrt{2}$	1.414 213 562 4	0.150 514 997 8
$\sqrt{3}$	1.732 050 807 6	0.238 560 627 4
$\sqrt{10}$	3.162 277 660 2	0.500 000 000 0
$\ln \pi$	1.144 729 885 8	0.058 703 021 2
$e^\gamma$	23.140 692 632 8	1.364 376 353 8
Euler constant $\gamma$	0.577 215 664 9	1.761 338 108 8
1 radian	57°29'577 9513 1	1.758 122 632 4
	3437746 770 78	3.536 273 882 8
	206 264°806 25	5.314 425 133 2
1°	0°017 453 292 5	2.241 877 367 6
1'	0°000 290 888 2	4.463 726 117 2
1"	0°000 004 848 1	6.685 574 866 8
Square degrees on a sphere = $129\ 600/\pi = 41\ 252.961\ 24$		
Square degrees in a steradian = $32\ 400/\pi^2 = 3282.806\ 35$		
For Gaussian distribution $(1/\sigma\sqrt{2\pi})\exp(-x^2/2\sigma^2)$		
Probable error/standard error = $r/\sigma = 0.674\ 489\ 750\ 2$		
Probable error/average error = $r/\eta = 0.845\ 347\ 539\ 4$		
$\sigma/\eta = 1.253\ 314\ 137$ , $\rho = (r/\sigma)/\sqrt{2} = 0.476\ 936\ 276\ 2$		




Figure A-1 Electromagnetic spectrum.<sup>1</sup>